

2001140570 MEDLINE
DOCUMENT NUMBER: 21067498 PubMed ID: 11150547
TITLE: Analysis of antigen presenting cell derived exosomes, based on immuno-magnetic isolation and flow cytometry.
AUTHOR: Clayton A; Court J; Navabi H; Adams M; Mason M D; Hobot J A; Newman G R; Jasani B
CORPORATE SOURCE: Section of Clinical Oncology, University of Wales College of Medicine, Velindre Hospital, Whitchurch, Cardiff CF14 2TL, UK.. aled.clayton@velindre-tr.wales.nhs.uk
SOURCE: JOURNAL OF IMMUNOLOGICAL METHODS, (2001 Jan 1) 247 (1-2) 163-74.
Journal code: 1305440. ISSN: 0022-1759.
PUB. COUNTRY: Netherlands
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200103
ENTRY DATE: Entered STN: 20010404
Last Updated on STN: 20010404
Entered Medline: 20010308

AB We present a simple yet powerful method for the isolation and analysis of exosomes released by antigen-presenting cells (APC). Exosomes are small **vesicles** (40-90 nm) released by APC, and may have an immuno-regulatory function in vivo. Such exosomes originate from MHC class II peptide loading compartments and, as such, express high levels of MHC Class II. We have utilised magnetic beads, coated with monoclonal antibodies specific for HLA DP, DQ, DR for the specific isolation of exosomes from cell-free supernatants. Beads coated with exosomes are subsequently stained with conjugated antibodies, and analysed by flow cytometry. Characterisation of exosomes by this method demonstrated that exosomes derived from B-lymphocytes express abundant MHC **Class I** and **Class II** molecules. Other immunologically important molecules detected included the co-stimulatory molecules B7.1 (CD80) and B7.2 (CD86). The adhesion molecule ICAM-1 (CD54) was also detected. These exosomes also expressed the **B cell** marker CD20, and the complement inhibitory protein CD59. The expression of CD63, a lysosomal marker, was variable, and there was no detectable expression of transferrin receptor (CD71). Monocyte derived dendritic cells (cultured for 7 days in GM-CSF/IL-4), demonstrated an immature phenotype, and secreted exosomes with a similar phenotype, with abundant MHC molecules. The expression of CD63 was consistently strong, and the MHC **Class I**-like molecule CD1a was also present, suggesting a possible function in the presentation of lipid antigens. Again CD59 was expressed suggesting a possible role for APC exosomes in complement regulation. There was no detectable CD71, CD40, CD14, CD20 or CD83. Modification of the extraction protocol allowed a comparative analysis of exosome secretion under various conditions. Treatment of cells with calcium ionophore, or phorbol ester resulted in apparent increases in exosome release, while the phosphatidyl inositol 3-kinase inhibitor, wortmannin, reduced exosome secretion. The immuno-magnetic isolation and analysis of exosomes is a versatile and rapid tool for the analysis of APC exosomes, and may prove a valuable tool for the study of exosome biology.